

522-1783

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN RE THE APPLICATION OF)
) Examiner:
Frank Luyten et al.)
) Group Art Unit No.
SERIAL NO.: To Be Assigned)
)
FILED: Herewith)
)
FOR: In Vivo Assay and Molecular Markers)
 for Testing the Phenotypic Stability of)
 Cell Population and Selected Cell)
 Population for Autologous Transplantation)

AMENDMENT ACCOMPANYING APPLICATION

Honorable Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Dear Sir:

The present application is the national filing of international application number PCT/BE00/00118. Before calculation of the national filing fee for the United States, it is requested that the application be amended as follows:

IN THE CLAIMS:

Cancel claims 1 through 28 (the only claims in the international application) without prejudice and substitute new claims 29 through 50 as follows:

29. An *in vivo* assay to identify molecular markers linked to phenotypic stability of a chondrocyte cell population comprising:

30. An assay to identify molecular markers according to claim 29, comprising using freshly isolated or serially passaged cells using differential gene expression analysis methods including differential display, subtractive hybridization, subtracted libraries or cDNA chips and cDNA arrays.

32. A method to identify cells having chondrocyte phenotypic stability according to claim
31 further comprising determining that activin-like kinase-1 (ALK-1) is not expressed, and/or
markers co-detectable with these markers and/or specific reporter constructs associated with
these markers are not expressed.

33. A method to identify cells with chondrocyte phenotypic stability comprising hybridising to messenger RNA from cells, sets of DNA probes provided on DNA arrays or DNA chips.
34. A method to identify phenotypically stable primary chondrocytes and chondrocytes, after at least one passage, that remained phenotypically stable comprising detecting sets of positive markers, said positive markers being selected from expressed BMP-2, FGFR-3, markers co-detectable with these markers or specific reporter constructs associated with these markers or markers determined by an in vivo assay according to claim 29.
35. Method to monitor passage by passage cell expansion and/or to predict when cell expansion must be stopped and/or to recover cells that have already lost their phenotypic stability only when needed and/or to provide a means for quality control of cells to be used for autologous cell transplantation and/or selecting from a cell population only those cells that retain their chondrocyte phenotypic stability, comprising detecting the expression of molecular markers of chondrocyte phenotypic stability selected from the group of:
 - markers determined by the assay comprising: a) injecting intramuscularly or subcutaneously in a non-human animal a suspension of isolated or expanded cells in an iso-osmotic liquid, the same suspension comprising articular chondrocytes in an amount equivalent to at least 1×10^6 chondrocytes as applied to immune-deficient mice, b) allowing the formation of cartilaginous tissue, c) sacrificing the animal, d) evaluate the formed cartilage histologically for stable, non-vascularised cartilage in vivo, and e) identify positive or negative molecular

–FGFR-3

pharmaceutical composition, that a treatment administered to the selected cells can hamper or enhance the anchorage-independent growth of said population as well as its phenotypic stability;

the in vivo assay comprising: subcutaneous or intramuscular injection in a non-human animal of a suspension of the selected cells in an iso-osmotic liquid, the same suspension comprising articular chondrocytes in an amount equivalent to at least 1×10^6 chondrocytes as applied to immune-deficient mice.

42. An *in vivo* assay

to predict the outcome of autologous cell transplantation using the cells selected by the method of claim 35

or to optimize cell culture conditions and manufacturing processes for a specific application in tissue engineering of cells selected by the method of claim 35,

or to evaluate the possibility that for the cells selected by the method of claim 35, said selected cells being used for autologous cell transplantation or being used in a pharmaceutical composition, that a treatment administered to the selected cells can hamper or enhance the anchorage-independent growth of said population as well as its phenotypic stability;

the in vivo assay comprising: subcutaneous or intramuscular injection in a non-human animal of a suspension of the selected cells in an iso-osmotic liquid, the same suspension

comprising articular chondrocytes in an amount equivalent to at least 1×10^6 chondrocytes as applied to immune-deficient mice.

43. Transplanting cells to a connective tissue site in a patient or seeding with cells any prosthetic device intended to be anchored into a mammal, said cells retaining their chondrocyte phenotypic stability and selected according to the method of claim 33.

44. Transplanting cells to a connective tissue site in a patient or seeding with cells any prosthetic device intended to be anchored into a mammal, said cells retaining their chondrocyte phenotypic stability and selected according to the method of claim 35.

45. A therapeutic composition for humans including cells selected according to claim 33, optionally further including at least a pharmaceutically acceptable carrier and/or a growth factor.

46. A therapeutic composition for humans including cells selected according to claim 35, optionally further including at least a pharmaceutically acceptable carrier and/or a growth factor.

47. A diagnostic comprising the DNA chips of claim 33.

48. A diagnostic for quality control of chondrocyte phenotypic stability. comprising at least one antibody selected from:

—an antibody specifically recognizing part of the extracellular domain of FGFR-3, obtained by immunizing animals with FGFR-3 or with a fragment thereof or with the synthetic peptide, TGLVPSEKVLVGPQRLQVLNASHEDSGAYSCRQRLTQRVL,

50. A cell culture exhibiting chondrocyte phenotypic stability in which the cells do not express activin-like kinase-1 (ALK-1) and/or markers co-detectable with this marker and/or specific reporter constructs associated with these markers as molecular markers negatively associated with chondrocyte phenotypic stability